



# Guide to Gamifant® (emapalumab-lzsg) Dosing and Administration

## Indication and Usage

Gamifant® (emapalumab-lzsg) is an interferon gamma (IFN $\gamma$ )-blocking antibody indicated for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy.

## Important Safety Information

Before initiating Gamifant, patients should be evaluated for infection, including latent tuberculosis (TB). Prophylaxis for TB should be administered to patients who are at risk for TB or known to have a positive purified protein derivative (PPD) test result or positive IFN $\gamma$  release assay.

**Please click [here](#) for Important Safety Information and full Prescribing Information.**

# Gamifant® (emapalumab-lzsg) Dosing and Administration

## Table of Contents

- 3** Important Safety Information
- 4** Primary HLH overview and importance of IFN $\gamma$
- 5** Gamifant clinical trial data
- 6** Supplies required for Gamifant preparation and infusion
- 6** Storing and handling of vials
- 6** How supplied
- 7** Premedication and concomitant medications
- 7** Infusion-related reactions
- 8** Calculate the Gamifant dose
- 9** Calculate the total infusion volume
- 9** Calculate the number of vials needed
- 10** Select the appropriate bags or syringes
- 11** Prepare Gamifant dilution
- 11** Storage of diluted solution
- 12** Parameters to consider when preparing infusion solution
- 13** Administering Gamifant by IV infusion
- 14** Titration intervals
- 14** Dose modification—response criteria for dose increase
- 15** Gamifant Patient Services
- 16** Ordering Gamifant

Please click here for [Important Safety Information](#)  
and full [Prescribing Information](#).



# Indication and Usage and Important Safety Information

## Indication and Usage

Gamifant® (emapalumab-lzsg) is an interferon gamma (IFN $\gamma$ )-blocking antibody indicated for the treatment of adult and pediatric (newborn and older) patients with primary hemophagocytic lymphohistiocytosis (HLH) with refractory, recurrent, or progressive disease or intolerance with conventional HLH therapy.

## Important Safety Information

Before initiating Gamifant, patients should be evaluated for infection, including latent tuberculosis (TB). Prophylaxis for TB should be administered to patients who are at risk for TB or known to have a positive purified protein derivative (PPD) test result or positive IFN $\gamma$  release assay.

During Gamifant treatment, patients should be monitored for TB, adenovirus, Epstein-Barr virus (EBV), and cytomegalovirus (CMV) every 2 weeks and as clinically indicated.

Patients should be administered prophylaxis for herpes zoster, *Pneumocystis jirovecii*, and fungal infections prior to Gamifant administration.

Do not administer live or live attenuated vaccines to patients receiving Gamifant and for at least 4 weeks after the last dose of Gamifant. The safety of immunization with live vaccines during or following Gamifant therapy has not been studied.

## Infusion-Related Reactions

Infusion-related reactions, including drug eruption, pyrexia, rash, erythema, and hyperhidrosis, were reported with Gamifant treatment in 27% of patients. In one-third of these patients, the infusion-related reaction occurred during the first infusion.

## Adverse Reactions

In the pivotal trial, the most commonly reported adverse reactions ( $\geq 10\%$ ) for Gamifant included infection (56%), hypertension (41%), infusion-related reactions (27%), pyrexia (24%), hypokalemia (15%), constipation (15%), rash (12%), abdominal pain (12%), CMV infection (12%), diarrhea (12%), lymphocytosis (12%), cough (12%), irritability (12%), tachycardia (12%), and tachypnea (12%).

Additional selected adverse reactions (all grades) that were reported in less than 10% of patients treated with Gamifant included vomiting, acute kidney injury, asthenia, bradycardia, dyspnea, gastrointestinal hemorrhage, epistaxis, and peripheral edema.

Please [click here](#) for full Prescribing Information.



# About primary HLH

Primary HLH is a genetic disorder that typically occurs in infancy and early childhood, manifesting mostly during the first year of life, but may also occur in teens and adults.<sup>1,2</sup>


**Primary HLH presents as a heterogeneous syndrome of rapidly progressive, life-threatening symptoms that can quickly become fatal unless diagnosed and treated. Common symptoms may include<sup>1,3,4</sup>:**



- Fever
- Infection
- Rash



- Hepatosplenomegaly
- Liver impairment
- Jaundiced appearance



- Hyperferritinemia
- Coagulation defects
- Severe cytopenia



- Seizure



- Pulmonary dysfunction

Massive overexpression of IFN $\gamma$  is central to the “cytokine storm,” the uncontrolled release of inflammatory cytokines and overactivation of phagocytes that give the syndrome its name<sup>5</sup>

## Important Safety Information

During Gamifant treatment, patients should be monitored for TB, adenovirus, Epstein-Barr virus (EBV), and cytomegalovirus (CMV) every 2 weeks and as clinically indicated.

Patients should be administered prophylaxis for herpes zoster, *Pneumocystis jirovecii*, and fungal infections prior to Gamifant administration.


Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Gamifant<sup>®</sup> (emapalumab-lzsg) is the first and only approved treatment for primary HLH<sup>6,7</sup>

**In clinical studies, Gamifant was shown to be effective treatment for primary HLH in patients with refractory, recurrent, or progressive disease or who were intolerant of conventional treatment.<sup>6</sup>**

- Multicenter, open-label, single-arm study of pharmacokinetics, efficacy, and safety of Gamifant in patients with suspected or confirmed primary HLH who had refractory, recurrent, or progressive disease during conventional HLH therapy or were intolerant to it<sup>6</sup>
  - **SAFETY** was evaluated in **34 patients**, 7 of whom were treatment naïve
  - **EFFICACY** was evaluated solely in **27 pediatric patients** who had already received conventional HLH treatment
- Patients had received a median of 3 prior agents as part of standard care before enrollment into the trial; prior regimens included combinations of dexamethasone, etoposide, cyclosporine A, and anti-thymocyte globulin<sup>6</sup>
- Median age of patients in the study was 1 year (range: 0.1-13 years)<sup>6</sup>



Primary endpoint was overall response rate (ORR) at the end of treatment, defined as achievement of either complete or partial response or HLH improvement.<sup>6</sup>

- **In patients with unsatisfactory response to conventional treatments, Gamifant achieved 63% ORR (95% CI: 0.42, 0.81; P = 0.013)<sup>6</sup>**



Secondary endpoints included time to response, durability of response, steroid reduction by 50% or more of baseline dose, and patients proceeding to hematopoietic stem cell transplantation (HSCT) when indicated.<sup>7</sup>

- **70% of patients (19/27) proceeded to HSCT<sup>6</sup>**

## Important Safety Information

Do not administer live or live attenuated vaccines to patients receiving Gamifant and for at least 4 weeks after the last dose of Gamifant. The safety of immunization with live vaccines during or following Gamifant therapy has not been studied.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Supplies required for Gamifant® (emapalumab-lzsg) preparation and infusion

Gamifant is administered as an intravenous infusion over 1 hour twice a week (every 3 to 4 days) until HSCT is performed. You will need the following supplies:

- Gamifant single-dose vials
- Gamma irradiated, latex-free, polyvinyl chloride (PVC)-free syringe (20 mL or 60 mL). Do not use with ethylene oxide-sterilized syringes
- Non-PVC polyolefin infusion bag (dependent on volume needed)
- 0.9% sodium chloride for injection, USP
- Intravenous line with sterile, non-pyrogenic, low-protein binding 0.2 µm in-line filter

## Storage and handling



- Store in a refrigerator at 2°C to 8°C (36°F to 46°F). Gamifant contains no preservative



- Store in original carton to protect from light



- DO NOT FREEZE OR SHAKE



- Do not transport via pneumatic tube

## How supplied

**Gamifant Injection is supplied in the following packaging configurations:**

- NDC 72171-501-01—containing one 10 mg/2 mL (5 mg/mL) single-dose vial
- NDC 66658-505-01—containing one 50 mg/10 mL (5 mg/mL) single-dose vial
- NDC 66658-510-01—containing one 100 mg/20 mL (5 mg/mL) single-dose vial



## Important Safety Information Infusion-Related Reactions

Infusion-related reactions, including drug eruption, pyrexia, rash, erythema, and hyperhidrosis, were reported with Gamifant treatment in 27% of patients. In one-third of these patients, the infusion-related reaction occurred during the first infusion.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Premedication and concomitant medications

## Premedication

- Administer prophylaxis for herpes zoster, *Pneumocystis jirovecii*, and for fungal infections prior to Gamifant® (emapalumab-lzsg) administration

## Concomitant medications

- Gamifant should be given concomitantly with dexamethasone
- For patients who are not receiving baseline dexamethasone treatment, begin dexamethasone at a daily dose of at least 5 to 10 mg/m<sup>2</sup> the day before Gamifant treatment begins
- Patients who are receiving baseline dexamethasone may continue their regular dose provided the dose is at least 5 mg/m<sup>2</sup>
- Dexamethasone can be tapered according to the judgment of the treating physician

## Infusion-related reactions

- Mild to moderate infusion-related reactions, including drug eruption, pyrexia, rash, erythema, and hyperhidrosis, were reported with Gamifant in 27% of patients
- In one-third of these patients, the infusion-related reactions occurred during the first infusion
- No infusion-related reactions led to premature withdrawal or were reported as serious adverse events

### Monitor patients for infusion-related reactions

Interrupt infusion for infusion-related reactions and institute appropriate medical management prior to continuing infusion at a slower rate

## Important Safety Information

### Adverse Reactions

In the pivotal trial, the most commonly reported adverse reactions (≥10%) for Gamifant included infection (56%), hypertension (41%), infusion-related reactions (27%), pyrexia (24%), hypokalemia (15%), constipation (15%), rash (12%), abdominal pain (12%), CMV infection (12%), diarrhea (12%), lymphocytosis (12%), cough (12%), irritability (12%), tachycardia (12%), and tachypnea (12%).

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Steps in preparing and administering Gamifant<sup>®</sup> (emapalumab-lzsg) to patients

## ► **STEP 1** Calculate the Gamifant dose


There are four variables to any Gamifant dose infusion

- Patient weight in kg
- Desired mg/kg dose
- Desired total infusion volume (to be administered over 1 hour)
- Patient condition (restrict total infusion volume; see chart on [page 9](#))



### **Record actual patient weight**

The weight of the patient must be taken prior to preparation of Gamifant for administration, ideally on the same day as the infusion.



### **Select the patient dose in mg**

(Can be 1 mg/kg, 3 mg/kg, 6 mg/kg, or 10 mg/kg)  
Patient weight (kg) x dose ([selected] mg/kg) = total mg of Gamifant needed.

## **Important Safety Information**

### **Adverse Reactions**

Additional selected adverse reactions (all grades) that were reported in less than 10% of patients treated with Gamifant included vomiting, acute kidney injury, asthenia, bradycardia, dyspnea, gastrointestinal hemorrhage, epistaxis, and peripheral edema.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).





## ► STEP 2 Calculate the total infusion volume and number of vials needed

### Recommended infusion volumes based on dose, infusion concentration, and patient weight

Patient Weight (kg)		3	5	10	15	20	25	30	40	50	60	70	90
Dose	Gamifant concentration in infusion solution												
1 mg/kg	0.5 mg/mL	6	10										
1 mg/kg	1 mg/mL			10	15	20	25	30	40	50	60	70	90
3 mg/kg	1 mg/mL	9	15	30	45	60	75	90					
3 mg/kg	2 mg/mL								60	75	90	105	135
6 mg/kg	2 mg/mL	9	15	30	45	60	75	90					
6 mg/kg	2.5 mg/mL								96	120	144	168	216
10 mg/kg	2 mg/mL	15	25	50	75								
10 mg/kg	2.5 mg/mL					80	100	120	160	200	240	280	360

Total mg of Gamifant® (emapalumab-lzsg) needed (from Step 1) ÷ selected mg/mL concentration = Total mL of Gamifant needed

**Note:** In cases where patient condition requires restriction of total infusion volume, higher concentration of infusion solution than those recommended can be used as long as the final concentration of infusion solution remains ≤ 2.5 mg/mL. Do not dilute product to < 0.25 mg/mL.

### Calculate the number of vials needed per dose of Gamifant Gamifant is available as 20 mL, 10 mL, or 2 mL vials

- Total mg of Gamifant needed ÷ 5 mg/mL Gamifant concentration
- Divide by 20 mL, 10 mL, or 2 mL for # of vials of Gamifant
- Vials are single-use only. Any remaining drug must be discarded

### Important Safety Information

Before initiating Gamifant, patients should be evaluated for infection, including latent tuberculosis (TB). Prophylaxis for TB should be administered to patients who are at risk for TB or known to have a positive purified protein derivative (PPD) test result or positive IFN $\gamma$  release assay.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



## Vial calculation example

If a patient weighs 5 kg and a 6 mg/kg dose is selected, then the total dose of Gamifant needed is 30 mg.

To calculate the number of vials needed for a 30 mg dose, divide by 5 mg/mL.

- 30 mg divided by 5 mg/mL = 6 mL of Gamifant
- 6 mL would require one 10 mL vial of Gamifant, which would produce 4 mL of waste. Or, select three 2 mL vials of Gamifant for exactly 6 mL of drug

## ► STEP 3 Select the appropriate bags or syringes

Depending on the dose to be administered and the weight of the patient, the diluted sterile concentrate can be administered either in 20 mL or 60 mL syringes or in a 0.9% sodium chloride for injection, USP infusion bag of the appropriate size, depending on the volume to be infused.

**For example, if the volume of the infusion solution (taking into account priming of the infusion line) is  $\leq 56$  mL, one syringe is to be used:**

- 20 mL syringe, when the volume to be delivered is  $\leq 16$  mL
- 60 mL syringe, when the volume to be delivered is  $> 16$  mL and  $\leq 56$  mL

**Note:** Do not use with ethylene oxide-sterilized syringes

**If  $> 56$  mL, an infusion bag is to be used:**

- 100 mL bag, when the volume to be delivered is  $\leq 80$  mL
- 250 mL bag, when the volume to be delivered is  $> 80$  mL and  $\leq 230$  mL
- 500 mL bag (if available) or 2 x 250 mL bags, when the volume to be delivered is  $> 230$  mL

## Important Safety Information

During Gamifant treatment, patients should be monitored for TB, adenovirus, Epstein-Barr virus (EBV), and cytomegalovirus (CMV) every 2 weeks and as clinically indicated.

Patients should be administered prophylaxis for herpes zoster, *Pneumocystis jirovecii*, and fungal infections prior to Gamifant administration.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).




## ► STEP 4 Prepare Gamifant® (emapalumab-lzsg) dilution

### Prepare the solution for infusion as follows:

- After removing from refrigerator, inspect Gamifant vials visually for particulate matter and discoloration prior to dilution. Gamifant is a clear to slightly opalescent, colorless to slightly yellow liquid. Do not administer if discolored or foreign particulate matter is present
- Withdraw the necessary amount of Gamifant solution and dilute with 0.9% Sodium Chloride Injection, USP to a maximum concentration of 2.5 mg/mL. Do not dilute product to less than 0.25 mg/mL
- Discard any unused portion left in the vial(s). Gamifant vials are for single use only
- The diluted solution can be placed in either a syringe or an infusion bag, depending on the volume needed



- Gently invert the infusion bag or syringe several times to ensure complete and homogeneous distribution of Gamifant



- DO NOT SHAKE

- Once the infusion solution is prepared, it should be clearly labeled for administration to the patient

Please see [page 12](#) for more information on infusion concentration parameters

### Storage of diluted solution

**Gamifant vials do not contain a preservative. If not administered immediately:**

- Store the diluted solution of Gamifant under refrigeration at 2°C to 8°C (36°F to 46°F) for no more than 4 hours from the time of dilution
- If refrigerated, allow the diluted solution to come to room temperature prior to administration
- Do not freeze or shake

### Important Safety Information

Do not administer live or live attenuated vaccines to patients receiving Gamifant and for at least 4 weeks after the last dose of Gamifant. The safety of immunization with live vaccines during or following Gamifant therapy has not been studied.

Please click [here](#) for **Important Safety Information** and full **Prescribing Information**.



# Parameters to consider for the preparation of Gamifant<sup>®</sup> (emapalumab-lzsg) solution for infusion

**An appropriate concentration of Gamifant in the infusion solution has to be chosen in order to optimize the final volume of the solution to be infused.**

- The maximum volume of non-diluted drug (5 mg/mL) to be administered should not exceed 50% of the total volume of the final solution. Therefore the maximum concentration to be used should be 2.5 mg/mL
- The total volume to be administered should take into account pediatric infusion guidelines:
  - For patients weighing less than 10 kg, the maximum volume to be administered should be 4 mL/kg/hr
  - For patients weighing between 10 and 20 kg, the maximum volume to be administered should be 6 mL/kg/hr

**The volume of the infusion solution to be prepared depends on priming or flushing.**

- **Priming**

The volume of the infusion line between the syringe and the intravenous catheter is taken into account in the final volume contained in the syringe or the bag.

- **Flushing**

The volume in the syringe or bag must be entirely delivered, which means that the infusion line is gently flushed with saline once the infusion is completed. If flushing is part of the infusion process to deliver the full dose, it should be performed immediately after the end of the infusion.

## Important Safety Information

### Infusion-Related Reactions

Infusion-related reactions, including drug eruption, pyrexia, rash, erythema, and hyperhidrosis, were reported with Gamifant treatment in 27% of patients. In one-third of these patients, the infusion-related reaction occurred during the first infusion.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



## ► **STEP 5 Administer Gamifant® (emapalumab-lzsg) by IV infusion**

**Gamifant is administered as an intravenous infusion over 1 hour twice a week (every 3 to 4 days) until HSCT is performed or unacceptable toxicity.**

Discontinue Gamifant when a patient no longer requires therapy for the treatment of HLH.

- Administer Gamifant diluted solution intravenously over 1 hour through an intravenous line containing a sterile, non-pyrogenic, low-protein binding 0.2 µm in-line filter
  - The duration of the infusion should be adapted when the volume to be infused is above the usual pediatric infusion recommendations (see table on [page 9](#))



- Do not store any unused portion of the infusion solution for reuse. Any unused product or waste material should be disposed of in accordance with local requirements

## **Important Safety Information**


### **Adverse Reactions**

In the pivotal trial, the most commonly reported adverse reactions ( $\geq 10\%$ ) for Gamifant included infection (56%), hypertension (41%), infusion-related reactions (27%), pyrexia (24%), hypokalemia (15%), constipation (15%), rash (12%), abdominal pain (12%), CMV infection (12%), diarrhea (12%), lymphocytosis (12%), cough (12%), irritability (12%), tachycardia (12%), and tachypnea (12%).

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Gamifant® (emapalumab-lzsg) dosing can be modified according to patient response



Gamifant offers flexibility of dosing and can be incrementally titrated upward or downward according to the clinician's assessment of patient response.

After the patient's clinical condition is stabilized, decrease the dose to the previous level to maintain clinical response until HSCT.

Dexamethasone can also be tapered according to the judgment of the treating physician. See [page 7](#) for more details.

## Dose increases should be based on both clinician assessment of unsatisfactory improvement in clinical condition, AND at least one of the following:

Platelet count	If baseline < 50,000/mm <sup>3</sup> and no improvement to > 50,000/mm <sup>3</sup>
	If baseline > 50,000/mm <sup>3</sup> and less than 30% improvement
	If baseline > 100,000/mm <sup>3</sup> , any decrease to < 100,000/mm <sup>3</sup>
Fever	Persistence or recurrence
Neutrophil count	If baseline < 500/mm <sup>3</sup> and no improvement to > 500/mm <sup>3</sup>
	If baseline > 500-1000/mm <sup>3</sup> and decrease to < 500/mm <sup>3</sup>
	If baseline 1000-1500/mm <sup>3</sup> and decrease to < 1000/mm <sup>3</sup>
Ferritin	If baseline ≥ 3000 ng/mL and < 20% decrease
	If baseline < 3000 ng/mL and any increase to > 3000 ng/mL
Splenomegaly	Any worsening
Coagulopathy (both D-dimer and fibrinogen must apply)	D-dimer: if abnormal at baseline and no improvement
	Fibrinogen (mg/dL): if baseline levels ≤ 100 mg/dL and no improvement or if baseline levels > 100 mg/dL and any decrease to < 100 mg/dL

## Important Safety Information

### Adverse Reactions

Additional selected adverse reactions (all grades) that were reported in less than 10% of patients treated with Gamifant included vomiting, acute kidney injury, asthenia, bradycardia, dyspnea, gastrointestinal hemorrhage, epistaxis, and peripheral edema.

Please [click here](#) for **Important Safety Information** and full **Prescribing Information**.



**Start treatment with Gamifant® (emapalumab-lzsg)**

## Gamifant Patient Services

### The first step to access is completing the Start Form

The Start Form enrolls your patients in Gamifant Patient Services and can be used to initiate insurance benefit verification, coordinate product orders, and assess a patient's eligibility for financial assistance.

Simply download and complete the Gamifant Start Form found at [Gamifant.com](http://Gamifant.com) and fax it to Gamifant Patient Services at **1-866-895-7204**.

- Be sure to include copies of your patient's insurance and pharmacy benefit cards
- Double-check that all required fields have been completed so as not to delay access

Once the Start Form has been received, a member of the Gamifant Patient Services team will contact you to facilitate the ordering process and help ensure that your patient receives their treatment as quickly as possible. If your order is urgent, please call 1-833-597-6530.

**Urgent order? Questions? We are here to help.**



**1-833-597-6530**

### Important Safety Information

Before initiating Gamifant, patients should be evaluated for infection, including latent tuberculosis (TB). Prophylaxis for TB should be administered to patients who are at risk for TB or known to have a positive purified protein derivative (PPD) test result or positive IFN $\gamma$  release assay.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).



# Ordering Gamifant® (emapalumab-lzsg)

Gamifant is available for order via our specialty pharmacy or through one of our two specialty distributors. Please note that the patient's insurance coverage may dictate how Gamifant should be ordered.

To better understand your patient's benefits please contact **Gamifant Patient Services** at 833.597.6530.

## Specialty pharmacy:

### Biologics

Phone: 800.850.4306

## Specialty distributors:

### McKesson Plasma and Biologics

Phone: 877.625.2566

Fax: 888.752.7626

Email: [mpborders@mckesson.com](mailto:mpborders@mckesson.com)

Online: [connect.mckesson.com](http://connect.mckesson.com)

### McKesson Specialty Health

Phone: 855.477.9800

Online: [mckessonspecialtyhealth.com](http://mckessonspecialtyhealth.com)

## Important Safety Information

During Gamifant treatment, patients should be monitored for TB, adenovirus, Epstein-Barr virus (EBV), and cytomegalovirus (CMV) every 2 weeks and as clinically indicated.

Patients should be administered prophylaxis for herpes zoster, *Pneumocystis jirovecii*, and fungal infections prior to Gamifant administration.

Please click here for [Important Safety Information](#) and full [Prescribing Information](#).





For more information, please visit:  
[Gamifant.com](http://Gamifant.com)

**References:** **1.** Jordan MB, Allen CE, Weitzman S, Filipovich AH, McClain KL. How I treat hemophagocytic lymphohistiocytosis. *Blood*. 2011;118(15):4041-4052. doi:10.1182/blood-2011-03-278127. **2.** Sepulveda FE, de Saint Basile G. Hemophagocytic syndrome: primary forms and predisposing conditions. *Curr Opin Immunol*. 2017;49:20-26. <http://dx.doi.org/10.1016/j.coi.2017.08.004>. **3.** Lehmsberg K, Nichols KE, Henter J-I, et al. Consensus recommendations for the diagnosis and management of hemophagocytic lymphohistiocytosis associated with malignancies. *Haematol*. 2015;100(8):997-1004. **4.** Marsh RA, Haddad E. How I treat primary haemophagocytic lymphohistiocytosis. *Br J Haematol*. 2018;182(2):185-199. doi:10.1111/bjh.15274. **5.** Price B, Lines J, Lewis D, Holland N. Haemophagocytic lymphohistiocytosis: a fulminant syndrome associated with multiorgan failure and high mortality that frequently masquerades as sepsis and shock. *S Afr Med J*. 2014;104(6):401-406. doi:7196/samj.7810. **6.** Gamifant [prescribing information]. Stockholm, Sweden: Biovitrum AB; 2020. **7.** Locatelli F, Jordan MB, Allen C, et al. Emapalumab in children with primary hemophagocytic lymphohistiocytosis. *N Engl J Med*. 2020;382(19):1811-1822.

Please click here for [Important Safety Information](#)  
and full [Prescribing Information](#).



Gamifant is a registered trademark, owned by Sobi AG and is marketed by Sobi, Inc.  
© 2020 Swedish Orphan Biovitrum. All rights reserved. PP-9261 09/20